



LYMPHA PRESS ORDER FORM

DATE OF ORDER _____

Medical Solutions Supplier is the exclusive provider of Lympha Press.

Patient Name _____ DOB _____ Phone _____

Extremities for Treatment

LEGS LT RT With Abdomen (Lympha Pants)

ARMS LT RT With Trunk (Comfy Sleeve)

Special Request _____

Treatment Pressures _____ Frequency _____ / Day _____ MIN Length of Necessity _____ Months (99=lifetime)

Default Settings 40mmHg, TID/BID, 60 min

Diagnosis

- Q82.0 HEREDITARY LYMPHEDEMA
- I89.0 SECONDARY LYMPHEDEMA DUE TO:
 - VENOUS INSUFFICIENCY
 - TUMOR(S)
 - SCARRING OF THE LYMPH CHANNELS FROM CELLULITIS AND/OR LYMPHANGITIS
 - CANCER
- I97.2 POST MASTECTOMY SYNDROME
- I87.2 CHRONIC VENOUS INSUFFICIENCY
 - VARICOSE VEIN WITH ULCER
 - VENOUS HYPERTENSION WITH ULCER

Prescribing Physician

Last _____ First _____
NPI# _____
Phone _____ Fax _____
Email _____
Physician Signature _____
In Clinic Contact _____
Date _____

Please Include with Order

- **FAX COVER SHEET**
with contact person name and phone/fax number.
- **PATIENT DEMOGRAPHICS**
including insurance information.
- **LAST 4+ WEEKS OF CHART NOTES (if available)**
Please send over the last few office visit notes preferably 4 weeks apart.

Documentation Examples

1) DIAGNOSIS

LYMPHEDEMA, identified as one of the following:

- Hereditary (Q82.0)
- Secondary (I89.0)
- Post mastectomy (I97.2)

CHRONIC VENOUS INSUFFICIENCY (I87.2)

with VENOUS STASIS ULCER(s) (I83.0 or I87.31)

2) SEVERITY, INCLUDING SPECIFIC EXAMPLES:

- Stages 2 or 3 lymphedema
- Ankle and calf measurements
- Hyperplasia
- Hyperkeratosis
- Papillomatosis cutis lymphostatica
- Hyperpigmentation
- Lymphorrhea
- Elephantiasis... etc.

3) CONSERVATIVE TREATMENTS (spanning 4 weeks)

1. Elevation
2. Exercise (ex. foot flexors, walking, etc.)
3. Compression wrapping or class 2 compression sleeve/stocking

4) SYMPTOMS PERSIST

Sample Dictations

“Patient presents with stage 2 lymphedema. Recommend regular elevation, exercise, and class 2 compression stockings (or wrapping system applied).”

After 4 weeks (28-42 days)

“Despite regular and compliant use of conservative treatments, including exercise, elevation, and class 2 compression stockings, the patient still presents with stage 2 lymphedema.”

Lympha Press Contraindications

Use of the Lympha Press® system is not recommended in the presence of one or more of the following conditions:

- Known or suspected deep vein thrombosis (DVT) or pulmonary embolism
- During the inflammatory phlebitis process
- Acute infection of the affected limb
- Decompensated cardiac failure or other ischemic vascular disease
- Severe arteriosclerosis or other ischemic vascular disease
- Any circumstance where increased venous and lymphatic return is undesirable
- The abdominal area should not be treated during pregnancy



lymphapress.com
Questions? Call 800-734-0422